

ABSTRACT OF THE DISCLOSURE

Method of identification and quantitative analysis of alcohol(s) in a sample by mass spectrometry using stable isotope labeled internal standard is provided. Said internal standard is prepared by reaction of an authentic sample of said alcohol with a stable isotope labeled reagent, and is added to a sample containing said alcohol. Said alcohol in said sample is then quantitatively converted to a chemical compound of identical structure, except the stable isotope atoms, as that of said internal standard using a non-labeled reagent. Said sample is then extracted and the extract is analyzed by mass spectrometry. Identification and quantification of said alcohol are made from a plot of ion ratio of said converted alcohol to said internal standard versus alcohol concentration.

REFERENCES

US patent documents

5,559,038	September 24, 1996	J. Fred Kolhouse
6,358,996	March 19, 2002	Michael S. Alexander

Other references

Mark M. Kushnir et al, "Comparison of four derivatizing reagents for 6-acetylmorphine GC-MS analysis", *Journal of Analytical Toxicology*, July/August 1999, page 262-269, vol. 23.

Michael L. Smith et al, "Forensic drug testing for opiates. VI. Urine testing for hydromorphone, hydrocodone, oxymorphone, and oxycodone with commercial opiate immunoassay and GC-MS", *Journal of Analytical Toxicology*, Jan/Feb 1995, page 18-26, vol. 19.

J. G. Guillot et al, "Determination of heroin, 6-acetylmorphine, and morphine in biological fluids using their propionyl derivatives with ion trap GC-MS", *Journal of Analytical Toxicology*, Mar/April 1997, page 127-133, vol. 21.